arXiv:1502.01267v1 [math.OA] 4 Feb 2015

CHARACTERIZATION OF CENTRAL ELEMENTS BY INEQUALITIES

AN.AN. NOVIKOV, O.E. TIKHONOV

Kazan Federal University, Kremlievskaia st. 18, 420008, Kazan, Russia
e-mail: a.hobukob@gmail.com, oleg.tikhonov@kpfu.ru

keywords: von Neumann algebra, C*-algebra, center of algebra, characterization, inequalities.
subclass: 46L05, 46L10, 47C15.

Partially supported by Russian Foundation for Basic Research, Grant 14-01-31358.

Abstract: We propose a list of inequalities which characterize central elements in von Neumann algebras and C*-algebras.

In this paper we study the possibility to distinguish central elements of operator algebras among all positive elements, by satisfying certain inequalities for positive functionals. A dual problem of characterizing tracial property by inequalities was considered, for instance, in [1, 2, 3, 4, 6, 8, 10, 12], and we apply some machinery of those papers.

Throughout the note, $\mathcal{M}$ stands for a von Neumann algebra, $\mathcal{M}^{sa}$, $\mathcal{M}^+$, and $\mathcal{M}^{\mathrm{pr}}$ denote the selfadjoint part, the positive part, and the set of all projections in $\mathcal{M}$, respectively. $\mathcal{Z}$ denotes the center of $\mathcal{M}$ and $\mathbf{1}$ denotes the identity operator. Let $\mathcal{M}_*$ denote the space of all normal functionals on $\mathcal{M}$, $\mathcal{M}_*^h$ and $\mathcal{M}_*^+$ denote its Hermitian and positive parts. We will use standard notation for multiplication of a functional φ by an operator x, namely, $x\varphi$, φx and $x\varphi x$ denote the linear functionals $y \mapsto \varphi(xy)$, $y \mapsto \varphi(yx)$ and $y \mapsto \varphi(xyx)$, respectively. Recall that a selfadjoint unitary operator in a Hilbert space is said to be a *symmetry*.

The proof of the following lemma is adapted from [10, Lemma 1].

Lemma 1. *Let* $a \in \mathcal{M}^+$. *If the inequality* $\varphi(sas) < \varphi(a)$ *holds for some symmetry* $s \in \mathcal{M}$ *and some positive normal functional* $\varphi \in \mathcal{M}_*^+$, *then there exist positive normal functionals* $\psi_1, \psi_2 \in \mathcal{M}_*^+$, *such that* $|\psi_1 - \psi_2|(a) > \psi_1(a) + \psi_2(a)$.

Proof. For $\lambda > 0$ define normal functionals $\psi_1^\lambda = \lambda s\varphi s + \varphi s + s\varphi + \lambda^{-1}\varphi$ and $\psi_2^\lambda = \lambda s\varphi s - \varphi s - s\varphi + \lambda^{-1}\varphi$. Since $\psi_1^\lambda = \lambda\, v_1^\lambda\, \varphi\, v_1^\lambda$ with $v_1^\lambda = s + \lambda^{-1}\mathbf{1}$ and $\psi_2^\lambda = \lambda\, v_2^\lambda\, \varphi\, v_2^\lambda$ with $v_2^\lambda = s - \lambda^{-1}\mathbf{1}$, those functionals are positive.

Let us demonstrate that $|\psi_1^\lambda - \psi_2^\lambda| = 2\varphi + 2s\varphi s$. Clearly, $|\psi_1^\lambda - \psi_2^\lambda| = 2|\varphi s + s\varphi|$. Observe that $\|\varphi s + s\varphi\| = (\varphi s + s\varphi)(s)$ since

$\|\varphi s + s\varphi\| \le \|\varphi s\| + \|s\varphi\| \le 2\|\varphi\| = 2\varphi(\mathbf{1})$ and $\|\varphi s + s\varphi\| \ge |(\varphi s + s\varphi)(s)| = (\varphi s + s\varphi)(s) = 2\varphi(\mathbf{1})$. By the construction of the absolute value of normal functional [11, the proof of Theorem III.4.2], we have $|\psi_1^\lambda - \psi_2^\lambda| = 2|\varphi s + s\varphi| = 2s(\varphi s + s\varphi) = 2(\varphi + s\varphi s)$.

With the notation $\varepsilon = (\varphi(a) - \varphi(sas))/\varphi(a)$, we obtain by a straightforward calculation $|\psi_1^\lambda - \psi_2^\lambda|(a) = 2(2-\varepsilon)\varphi(a)$ as well as $(\psi_1^\lambda + \psi_2^\lambda)(a) = (\lambda(1-\varepsilon) + \lambda^{-1})\varphi(a)$. Since $\min\limits_{\lambda>0}\{\lambda(1-\varepsilon) + \lambda^{-1}\} = 2\sqrt{1-\varepsilon}$ and $2-\varepsilon > \sqrt{1-\varepsilon}$ for any $\varepsilon \in (0,1)$, there exists $\lambda_0 > 0$ such that $2(2-\varepsilon) > \lambda_0(1-\varepsilon) + \lambda_0^{-1}$. Therefore $|\psi_1^{\lambda_0} - \psi_2^{\lambda_0}|(a) > (\psi_1^{\lambda_0} + \psi_2^{\lambda_0})(a)$. $\square$

Theorem 1. *For $a \in \mathcal{M}^+$ the following conditions are equivalent:*

- (i) *a lies in the center $\mathcal{Z}$ of $\mathcal{M}$;*
- (ii) *$pap \le a$ for each $p \in \mathcal{M}^{pr}$;*
- (iii) *$\varphi^+(a) \le \varphi_1(a)$ for each $\varphi \in \mathcal{M}_*^h$ and any decomposition $\varphi = \varphi_1 - \varphi_2$ with $\varphi_1, \varphi_2 \in \mathcal{M}_*^+$;*
- (iv) *the mapping $\varphi \mapsto \varphi^+(a)$ $(\varphi \in \mathcal{M}_*^h)$ is monotone, i. e., $\varphi, \psi \in \mathcal{M}_*^h$, $\varphi \le \psi$ imply $\varphi^+(a) \le \psi^+(a)$;*
- (v) *the mapping $\varphi \mapsto \varphi^+(a)$ $(\varphi \in \mathcal{M}_*^h)$ is subadditive, i. e., $(\varphi + \psi)^+(a) \le \varphi^+(a) + \psi^+(a)$ for all $\varphi, \psi \in \mathcal{M}_*^h$;*
- (vi) *the mapping $\varphi \mapsto |\varphi|(a)$ $(\varphi \in \mathcal{M}_*^h)$ is subadditive, i. e., $|\varphi + \psi|(a) \le |\varphi|(a) + |\psi|(a)$ for all $\varphi, \psi \in \mathcal{M}_*^h$.*

Proof. $(i) \Rightarrow (ii)$ For $a \in \mathcal{Z}^+$ and $p \in \mathcal{M}^{\mathrm{pr}}$, one has $a = pap + (\mathbf{1} - p)a(\mathbf{1} - p) \ge pap$.

$(ii) \Rightarrow (iii)$ For $\varphi \in \mathcal{M}_*^h$, let p be the support projection of φ^+ (see [11, Section III.4]). Then $\varphi^+(a) = (p\varphi p)(a) \le (p(\varphi + \varphi_2)p)(a) = (p\varphi_1 p)(a) = \varphi_1(pap) \le \varphi_1(a)$.

$(iii) \Rightarrow (iv)$ This follows from the equality $\varphi = \psi^+ - (\psi^- + (\psi - \varphi))$.

$(iv) \Rightarrow (v)$ For $\varphi, \psi \in \mathcal{M}_*^h$, one has $\varphi \le \varphi^+$ and $\psi \le \psi^+$, hence $\varphi + \psi \le \varphi^+ + \psi^+$. Then the condition (iv) entails $(\varphi + \psi)^+(a) \le \varphi^+(a) + \psi^+(a)$.

$(v) \Rightarrow (vi)$ By the condition (v), for $\varphi, \psi \in \mathcal{M}_*^h$, it holds $(\varphi + \psi)^+(a) \le \varphi^+(a) + \psi^+(a)$. Also, $(\varphi + \psi)^-(a) = (-\varphi - \psi)^+(a) \le (-\varphi)^+(a) + (-\psi)^+(a) = \varphi^-(a) + \psi^-(a)$. Hence $|\varphi + \psi|(a) = (\varphi + \psi)^+(a) + (\varphi + \psi)^-(a) \le \varphi^+(a) + \psi^+(a) + \varphi^-(a) + \psi^-(a) = |\varphi|(a) + |\psi|(a)$.

$(vi) \Rightarrow (i)$ If an operator $a \in \mathcal{M}^+$ satisfies (vi), then it follows from Lemma 1 that $\varphi(sas) = \varphi(a)$ for each symmetry s in $\mathcal{M}$ and any φ in $\mathcal{M}_*^h$. Therefore $a = sas$ for each symmetry s in $\mathcal{M}$. As it is easily seen, the latter implies that a commutes with each projection in $\mathcal{M}$ and therefore lies in $\mathcal{Z}$. $\square$

Remark 1. Clearly, for $a \in \mathcal{M}^+$, (v) is equivalent to the condition (*vii*) *the mapping* $\varphi \mapsto \varphi^+(a)$ $(\varphi \in \mathcal{M}_*^h)$ *is convex.*

As well, (*vi*) is equivalent to the conditions

(*viii*) *the mapping* $\varphi \mapsto |\varphi|(a)$ $(\varphi \in \mathcal{M}_*^h)$ *is convex.*

It follows from Theorem 1 that for $a \in \mathcal{M}^+$ each of the conditions (*vii*), (*viii*) is equivalent to $a \in \mathcal{Z}$.

Corollary 1. *For $a \in \mathcal{M}^+$, each of the conditions (i) – (viii) is equivalent to each of the following conditions:*

(*ix*) $|\varphi|(a) = \|a^{\frac{1}{2}}\varphi a^{\frac{1}{2}}\|$ *for all φ in $\mathcal{M}_*$;*

(*x*) *the mapping $\varphi \mapsto |\varphi|(a)$ is subadditive on $\mathcal{M}_*$,*
i. e., $|\varphi + \psi|(a) \le |\varphi|(a) + |\psi|(a)$ for all $\varphi, \psi \in \mathcal{M}_*$.

Proof. Let $a \in \mathcal{Z}^+$. For $\varphi \in \mathcal{M}_*$, let $\varphi = u|\varphi|$ be the polar decomposition [11, Section III.4]. Then $|\varphi|(a) = (u^*\varphi)(a) = \varphi(u^*a) = \varphi(a^{\frac{1}{2}}u^*a^{\frac{1}{2}}) = (a^{\frac{1}{2}}\varphi a^{\frac{1}{2}})(u^*) \le \|a^{\frac{1}{2}}\varphi a^{\frac{1}{2}}\|$. On the other hand, $\|a^{\frac{1}{2}}\varphi a^{\frac{1}{2}}\| = \|a^{\frac{1}{2}}u|\varphi|a^{\frac{1}{2}}\| = \|ua^{\frac{1}{2}}|\varphi|a^{\frac{1}{2}}\| \le \|a^{\frac{1}{2}}|\varphi|a^{\frac{1}{2}}\| = |\varphi|(a)$. Hence (*ix*) is satisfied.

(*ix*) $\Rightarrow$ (*x*). If (*ix*) holds true, then we have $|\varphi + \psi|(a) = \|a^{\frac{1}{2}}(\varphi + \psi)a^{\frac{1}{2}}\| \le \|a^{\frac{1}{2}}\varphi a^{\frac{1}{2}}\| + \|a^{\frac{1}{2}}\psi a^{\frac{1}{2}}\| = |\varphi|(a) + |\psi|(a)$.

Of course, (*x*) implies (*vi*). □

Remark 2. From the preceding proof, it is seen that we can add the following condition to the list (*i*) – (*x*) of equivalent conditions.

(*xi*) $|\varphi|(a) = \|a^{\frac{1}{2}}\varphi a^{\frac{1}{2}}\|$ *for all φ in $\mathcal{M}_*^h$.*

The following theorem is an analog of Gardner's characterization of traces by "triangle inequality" [6] (see, also, [8]).

Theorem 2. *A positive element a of $\mathcal{M}$ belongs to the center of $\mathcal{M}$ if and only if the inequality $|\varphi(a)| \le |\varphi|(a)$ holds for any φ in $\mathcal{M}_*$.*

Proof. Taking into account Corollary 1, for $a \in \mathcal{Z}$ and $\varphi \in \mathcal{M}_*$, we get $|\varphi(a)| = |(a^{\frac{1}{2}}\varphi a^{\frac{1}{2}})(\mathbf{1})| \le \|a^{\frac{1}{2}}\varphi a^{\frac{1}{2}}\| = |\varphi|(a)$.

Let $|\varphi(a)| \le |\varphi|(a)$ for any φ in $\mathcal{M}_*$, then $|(u\psi)(a)| = |\psi(ua)| \le \psi(a)$ for any unitary u and any positive normal ψ. Since the unit ball of $\mathcal{M}$ is the closed convex hull of the set of unitaries [9] (see also [7, 1.1.12]), it follows that $|\psi(xa)| \le \|x\|\psi(a)$ for every x in $\mathcal{M}$. But then the functional ψa attains its norm at $\mathbf{1}$ and is therefore positive [5, 2.1.9]. If $x \in \mathcal{M}^{sa}$ then $\psi(xa) = \overline{\psi(xa)} = \psi(ax)$. It follows that $xa = ax$ for any $x \in \mathcal{M}^{sa}$ and therefore a belongs to the center. □

Corollary 2. *A positive element a of a C^*-algebra $\mathcal{A}$ belongs to the center of $\mathcal{A}$ if and only if the inequality $|\varphi(a)| \le |\varphi|(a)$ holds for any φ in $\mathcal{A}^*$.*

Proof. Let $(\pi, \mathfrak{H})$ be the universal representation of $\mathcal{A}$ and $\mathcal{M}(\pi)$ be the universal enveloping von Neumann algebra [11, Section III.2]. By construction of $\mathcal{M}(\pi)$, the spaces $\mathcal{A}^*$ and $\mathcal{M}(\pi)_*$ are isometrically isomorphic in a natural way. For $\varphi \in \mathcal{A}^*$, we will denote by $\widetilde{\varphi}$ the corresponding functional in $\mathcal{M}(\pi)_*$. Note that $\widetilde{|\varphi|} = |\widetilde{\varphi}|$ by construction of absolute value [5, 12.2.7, 12.2.8].

Let a be a positive element of the center of $\mathcal{A}$. It is easy to see that $\pi(a)$ belongs to the center of $\mathcal{M}(\pi)$. Then for any $\varphi \in \mathcal{A}^*$ we have $|\varphi(a)| = |\widetilde{\varphi}(\pi(a))| \leq |\widetilde{\varphi}|(\pi(a)) = \widetilde{|\varphi|}(\pi(a)) = |\varphi|(a)$.

On the other hand, let a be a positive element of $\mathcal{A}$ and the inequality $|\varphi(a)| \leq |\varphi|(a)$ hold for any φ in $\mathcal{A}^*$. Then $|\widetilde{\varphi}(\pi(a))| \leq |\widetilde{\varphi}|(\pi(a))$ for any $\widetilde{\varphi}$ in $\mathcal{M}(\pi)_*$, which implies that $\pi(a)$ lies in the center of $\mathcal{M}(\pi)$, hence a lies in the center of $\mathcal{A}$. $\square$

Remark 3. One can easily adapt the conditions (iii) – (xi) to the case of C*-algebras and see that each of those modified conditions characterizes central elements.

REFERENCES

[1] A.M. Bikchentaev, Dokl. Math., **87** (1), 79-82 (2013).

[2] A.M. Bikchentaev, Sib. Math. J., **51** (6), 971-977 (2010).

[3] A.M. Bikchentaev, Math. Notes, **89** (3-4), 461-471 (2011).

[4] Dinh Trung Hoa, O.E. Tikhonov, Math. Notes, **88** (2), 177-182 (2010).

[5] J. Dixmier, *C*-algebras*, (North Holland Publishing co., Amsterdam, 1977).

[6] L.T. Gardner, Can. J. Math. **31**, 1322-1328 (1979).

[7] G. K. Pedersen, *C*-algebras and their automorphism groups*, (London Math. Soc., Monographs **14**, Academic Press, London/New York, 1979).

[8] G.K. Pedersen, E. Størmer, Can. J. Math. **34**, 370-373 (1982).

[9] B. Russo, H.A. Dye, Duke Math. J. **33**, 413-416 (1966).

[10] A.I. Stolyarov, O.E. Tikhonov, A.N. Sherstnev, Math.Notes **72** (3), 411-416 (2002).

[11] M. Takesaki, *Theory of operator algebras I*, (Encyclopaedia of Mathematical Sciences **124**, Reprint of the first (1979) edition, Operator Algebras and Non-commutative Geometry, 5 Springer-Verlag, Berlin, 2002).

[12] O.E. Tikhonov, Positivity, **9** (2), 259-264 (2005).